



Reg. No. : .....

Name : .....



Fourth Semester B.Tech. Degree Examination, May 2014  
(2008 Scheme)

08.401 : ENGINEERING MATHEMATICS – III (CMPUNERFHB)

Time: 3 Hours

Max. Marks: 100

**Instructions:** Answer **all** questions from Part – **A** each question carries 4 marks and **one full** question from **each** module of Part – **B** each full question carries 20 marks.

PART – A

1. Prove that  $f(z) = e^z$  is differentiable everywhere and find its derivative.
2. Prove that an analytic function with constant argument is constant.
3. Choose 'a' so that the function  $u = x^3 + axy^2$  is harmonic, find its harmonic conjugate.
4. Prove that a bilinear transformation preserves cross ratio.
5. Evaluate  $\int_0^{1+i} (x^2 - iy) dz$  along  $y = x$  and  $y = x^2$ . Are they equal ?
6. Using Cauchy's integral formula, evaluate  $\int_C \frac{e^{2z}}{(z+1)^4} dz$  where  $C$  is  $|z| = 2$ .
7. Obtain the Taylor series expansion of  $f(z) = \frac{1}{z^2}$  about  $z = 2$ .
8. Explain Newton-Raphson method.



9. Apply Lagrange's interpolation formula to find  $f(6)$  given  $f(1) = 2$ ,  $f(2) = 4$ ,  $f(3) = 8$ ,  $f(4) = 16$  and  $f(7) = 128$ .
10. The following table gives the values of  $f(x)$  at equal intervals of  $x$ .

$x :$	0	0.5	1	1.5	2	2.5	3
$f(x) :$	0	0.7071	1	1.2247	1.4142	1.5811	1.732

Evaluate  $\int_0^3 f(x) dx$  by

i) Simpson's  $\frac{1}{3}$  rule.

ii) Trapezoidal rule.

(10×4=40 Marks)

**PART - B**  
**Module - I**

11. a) Show that  $f(z) = \frac{xy^2(x + iy)}{x^2 + y^4}$  for  $z \neq 0$   
 $= 0$  for  $z = 0$   
 is not differentiable at  $z = 0$ .
- b) If  $f(z) = u + iv$  is analytic, prove that the families of curves  $u = c_1$  and  $v = c_2$  where  $c_1$  and  $c_2$  are constant cut orthogonally.
- c) Find the bilinear transformation that maps the points  $z = 1, i, -1$  onto  $w = i, 0, -i$ . Hence find the image of  $|z| < 1$ .
12. a) Prove that  $f(z) = xy + iy$  is everywhere continuous but not analytic.
- b) Given  $u + v = \frac{2 \sin 2x}{e^{2y} + e^{-2y} - 2 \cos 2x}$  find  $f(z) = u + iv$  by Milne-Thompson method.
- c) Determine the region of the  $w$ -plane into which the triangular region bounded by  $x = 1, y = 1$  and  $x + y = 1$  is mapped by  $w = z^2$ .



**Module – II**

13. a) Evaluate  $\int_C \frac{\sin \pi z^2 + \cos \pi z^2}{(z-1)^2(z-2)} dz$  where C is  $|z|=2$ .
- b) Obtain the Laurent's series expansion of  $f(z) = \frac{1}{(z-1)(z-2)}$  in  $0 < |z-1| < 1$  and hence find its residue about  $z=1$ .
- c) Evaluate  $\int_0^\infty \frac{1}{(a^2+x^2)^2} dx$  using Cauchy's residue theorem.

14. a) Determine the nature and singularities of

i)  $\frac{e^{2z}}{(z-1)^4}$

ii)  $ze^{\frac{1}{z^2}}$



b) Evaluate  $\int_{|z|=3} \frac{e^z}{(z+2)(z+1)^2} dz$  by Cauchy's residue theorem.

c) Evaluate  $\int_0^{2\pi} \frac{\cos 2\theta}{5+4\cos \theta} d\theta$ .

**Module – III**

15. a) Find the approximate value correct to three places of decimals of the real root which lies between  $-2$  and  $-3$  of the equation  $x^3 - 3x + 4 = 0$  by regula falsi method.
- b) Solve by Gauss-Seidal method given
- $$10x - 2y - z - 10 = 3$$
- $$-2x + 10y - z - w = 15$$
- $$-x - y + 10z - 2w = 27$$
- $$-x - y - 2z + 10w = -9.$$



c) From the following table :

$x :$	0.1	0.2	0.3	0.4	0.5	0.6
$f(x) :$	2.68	3.04	3.38	3.68	3.96	4.21

Find  $f(0.7)$  by appropriate interpolation formula.

16. a) Find the real root of  $x^3 - 2x = 5$  correct to three decimal places by bisection method.
- b) Find the value of  $y$  when  $x = 0.1, 0.2$  and  $0.3$  by Runge-Kutta method given  $y' = xy + y^2; y(0) = 1$ .
- c) Using Taylor series method solve  $y' = x^2 - y; y(0) = 1$  at  $x = 0.1, 0.2$  and  $0.3$ .

(3×20=60 Marks)

---